

Hygienic profile and starch content of traditional yogurts in north of Iran

Morteza Rahbar-Taramsari¹, Marjan Mahdavi-Roshan^{2*}, Kasra Hemmati³, Mojtaba Hedayati⁴

1- Department of Forensic Medicine, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran.

2- Cardiovascular Diseases Research Center, Department of Cardiology, Heshmat Hospital, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran.

3- School of Medicine, Guilan University of Medical Sciences, Rasht, Iran.

4- Department of Microbiology, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran.

This paper is open access under [Creative Commons Attribution-NonCommercial 4.0 International](https://creativecommons.org/licenses/by-nc/4.0/) license.



Submission: 9 March 2021

Revision: 18 April 2021

Acceptance: 18 April 2021

Abstract

Background and objective: Traditional dairies have been produced and consumed since long time ago. However, their quality status is not routinely examined. Therefore, we determined microbial content and starch level of traditional yogurts produced in north of Iran.

Materials and methods: Detection of Coliforms and *Escherichia coli* (by MPN method), *Staphylococcus aureus* (by surface plating on Baird Parker agar) and yeast/mold (by surface plating on Dichloran Rose-Bengal Chloramphenicol Agar) were conducted. Starch inclusion was determined by addition of iodine reagent to the products and monitoring of color change.

Results and conclusion: All samples were approved in Coliforms, *E. coli*, and *S. aureus* tests. Mold was detected in five samples and the all showed yeast growth more than accepted level. According to Iranian regulation about fermented foods containing yeast as starter that exempted from maximum limit of yeast, majority of yogurts were appropriate for consumption in view of yeast contamination. Most of the samples did not contain starch except for two yogurts containing high starch. Low microbial contamination and starch content in the samples was due to the use of safe raw materials and prominence of healthy microbes present in traditional dairies in Iran.

Keywords: Microbial contamination, starch, traditional yogurt, Iran

1. Introduction

Dairy industry is one of the most critical and potent commerce in the world. Consumption of milk-based products per day is recommended by dietitians all the time due to their health benefits in supplying the calcium need for body that is 1000 mg per 2000 kcal energy intake per day. Calcium content of one cup of milk is accounted

as 300 mg, of which approximately 32% is absorbed. Therefore, an average of three cups of milk or its equivalent in other products is required to supply the body's calcium need [1]. Among dairy products, several positive impacts are addressed to those underwent fermentation such as yogurt. For example, they are well tolerated by lactase deficient consumers because lactose (that

* Correspondence to: Marjan Mahdavi-Roshan, e-mail: mahdaviroshan@gums.ac.ir; Tel.: +98-13-32135352

is a prevalent disaccharide in milk) is fermented by indigenous or added starter species to lactic acid and other compounds [2]. Moreover, inclusion of healthy bacteria such as *Streptococcus thermophilus* and bifidobacteria through fermentation in dairy products are in favor of supplying the required folate to consumers [3].

Despite industrial production by addition of commercial starter bacteria or yeasts, traditional dairies are produced locally by people in rural or urban area by using sour dough. These products are popular in Asian countries. In Iran, many people tend to consume traditional dairies because of their natural ingredients, desired sharp flavor, and health benefits. Indeed, traditional yogurt is produced from milk and starter cultures and no processing-aid are added. Various probiotics such as *Lactobacillus*, *Streptococcus*, and *Leuconostoc* have been isolated from traditional yogurts, which is comparable to industrial yogurts containing two or three species of starter. This is due to the vast geographic area with natural sources in the Middle East countries that favors derivation of several microorganisms in traditional products [4]. In agreement, Tajabadi Ebrahimi et al. isolated three species of *Lactobacillus* from traditional cheese and yogurt in Iran. They characterized the isolates as probiotic with regard to acid and bile tolerance, pathogen antagonism, and cholesterol assimilation [5]. Similar results were further reported by Faghfoori et al. about probiotic *Lactobacillus* species isolated from traditional cheeses and yogurts prepared in north-west of Iran [6]. Other than *Lactobacillus*, *Pediococcus acidilactici* was isolated from traditional yogurts in Iran by Sharifi Yazdi et al. that showed probiotic properties [7]. Owing to inclusion of diverse viable cells, traditional yogurts might have more positive impact on the consumers than conventional yogurts through production of several bioactive compounds by fermentation of milk nutrients. It has been reported that starter microbes are able to produce bioactive peptides through fermentation of milk proteins, which pose health benefits in lowering serum cholesterol, blood sugar control, immune regulation, and antioxidant status [2,4]. Gamma amino butyric acid (GABA) is a bioactive non-protein amino acid and differentiated from glutamic acid is able to regulate sleep disorder, lower blood pressure, stimulate immune

response, and suppress stress. In the study carried out on probiotic bacteria isolated from traditional dairies in Iran, high potential of GABA production was reported for *Lactobacillus paracasei*, *L. plantarum*, and *P. acidilactici* [8]. Moreover, Hamzehlou et al. in 2018 characterized eleven *Lactobacillus* isolated from traditional yogurts produced in different regions of Iran. The bacteria had high potential in production of B-group vitamins and were suggested to be used in production of other functional foods [9]. Selenium is a vital micronutrient mainly found in marine foods and meat and can be delivered by supplements. However, its high intake can lead to some diseases such as diabetes, prostate cancer, and neurodegenerative diseases by species-dependent manner. Indeed, inorganic and some organic derivatives have higher toxic impact than the others [10]. Interestingly, biotransformation of selenite derivatives to its reduced elemental form by lower toxicity was reported for *L. brevis* LSe isolated from traditional sour yogurt in Iran. The authors found that the bacterium could grow in the presence of selenite ions and suggested that it can alleviate side effects of selenium overload in body [11]. Angiotensin I-converting enzyme inhibitory action was determined for bioactive peptides produced by active bacteria in traditional yogurts, through which an antihypertensive property is expected [12].

The approved health benefits of indigenous microorganisms derived from traditional yogurts have resulted in a great tendency to their consumption in recent decades. On the other hand, the consumers are not aware of hygienic status or unknown additives of traditional yogurts and it may lead to unexpected adverse outcomes especially in sensitive people. Therefore, the aim of current study was determination of microbial profile (*Coliform*, *Escherichia coli*, *Staphylococcus aureus*, yeast and mold) and starch content of traditional yogurts produced in north of Iran. At the end of research, we could conclude that whether the products can help people to intake functional ingredients or might put the consumers at risk of unexpected diseases.

2. Materials and methods

2-1- Materials

Sterile glass and plastic wares, wire loops, and inoculating needles were used during experi-

ments. The glass wares were washed and rinsed with distilled water, sterilized, and underwent endotoxin removal at 400 °C for 3 h. The culture media were accurately weighed and dissolved in distilled water according to the manufacturer's direction and were stored in refrigerator until use. The culture media of Lauryl Sulfate Tryptose broth, EC broth, and Tryptone Water and Indole reagent (mixture of isoamyl alcohol, para-dimethylaminobenzaldehyde, and concentrated hydrochloric acid) were used for Coliform and *E. coli* enumeration. The selective medium of Baird Parker agar was used for *S. aureus* determination. Yeast and mold determination was conducted on Dichloran Rose-Bengal Chloramphenicol agar. Iodine and potassium iodide were used for preparation of iodine reagent for starch determination test. All the media and chemical reagents were purchased from Merck Company (Germany).

2-2- Sample size and sampling procedure

Samples were collected from local markets in north or Iran (Rasht city). The following formula was used for detection of sample size.

$$n = \frac{z^2 \times \delta^2}{d^2}$$

Where, "n" is sample size, "z" is 1.96 for 95% confidence interval, "δ" is standard deviation (average of 0.16 was used based on the results of previous study [13]), and "d" is acceptable error (a factor between 0.01 and 0.1; the average of 0.066 was used). Therefore, a sample size of 23 was achieved. In practice, the city was divided to north, south, west, east, and center and 23 samples were collected randomly. The collected samples were labeled numerically (1-23), and then were transported to the microbiology lab at low temperature by a portable cool chamber.

2-3- Microbial analysis

All experiments were conducted under sterile condition to avoid cross-contamination. Positive control of bacterial and fungal isolates were prepared and used for comparison. For bacterial enumeration, serial dilutions were prepared firstly by adding 1 g of homogenized sample to 9 ml sterile saline solution (0.85% w/v in distilled water). Then, 1 ml of the first dilution was added to 9 ml of the dilutant in series up to reach the dilution of 10⁻⁶ [14]. Coliform/*E. coli*, coagulase

positive staphylococcus, and yeast/mold were enumerated by ISO 11866-1 | IDF 170-1, ISO 6888-1, and ISO 21527-1, respectively [15-17]. Incubation conditions were 37 °C for 24-48 h for the bacteria and 25 °C for five days for yeast/mold. The colonies grown after incubation were counted and expressed as colony forming unit per gram of the samples. In the next step, bacterial isolates were characterized by morphological and biochemical tests including IMViC series, evaluation of color and shape of colonies, gram staining, monitoring of motility, and ability to produce oxidase, catalase, urease, coagulase, and sulfide.

2-4- Starch determination

Approximately, 1-2 ml of iodine reagent (2.6 g iodine and 3 g potassium iodide in 200 ml of distilled water) was dropped into 5 ml of sample in test tube. The mixture was boiled and then left to cool to ambient temperature. Disappearance of blue color in the boiling sample and its further appearance in the cooled sample was indicator of starch in yogurt [18].

3. Results and discussion

Iran is a vast country with several natural foods assessed for their functional and health properties. Endemic traditional foods are of concern sometimes because they do not undergo hygienic processes that industrial products do. Traditional products are not monitored regularly and not-allowed additives may be added by producers. Codex alimentarius has classified yogurt as fermented milk and defined it as "a milk product obtained by fermentation of milk, by the action of suitable microorganisms and resulting in reduction of pH with or without coagulation (isoelectric precipitation). These starter microorganisms shall be viable, active, and abundant in the product to the date of minimum durability" [19]. Several adulterations are addressed in dairies to make the products cheaper, more palatable, and resistant to microbial contamination but they threaten the consumers' health in some cases [20]. They include addition of water, vegetable or animal protein, non-dairy fat, urea, melamine, starch, ammonium sulfate, formalin, and non-declared species of milk to the final products [21-24]. These can be fully detected by instrumental and complex analytical methods such as Raman

spectroscopy, high performance liquid chromatography-mass spectroscopy, and chemometrics approaches [23-25].

Because of people's concern about adverse effects of the non-permitted additives that are not detected visually in industrial products, some of them prefer to use traditional dairy products. In this regard, evaluation of microbial load as health index and starch level that is harmless but may cover technical faults were done in traditional yogurts collected from local markets in north of Iran.

3-1- Microbial load

Traditional dairies do not usually expose to pasteurization and may contain unacceptable level of pathogens. Although, it is supposed that the harmful microbes are destroyed in the products under fermentation process in the presence of healthy starters. A descriptive list of potential bacteria, yeasts, and molds in fermentation of dairies, beverages, cereals, coffee, cocoa, fishes,

fruits, meats, sourdough, soy, tea, vegetables, vinegar, and wine was reported by European Food and Feed Cultures Association [26].

According to Table 1, no growth of *E. coli* and acceptable number of coliforms were detected in all samples. Interestingly, no *S. aureus* was detected in the samples. Other than good hygienic status of the products, it might be due to the metabolism of healthy bacteria that led to pathogens' suppression, if existed. In corroboration, inhibitory effect of lactic acid bacteria active in fermented milks was reported by Gomez-Gallego et al. against *E. coli* and *Salmonella enteritidis* [3]. Cutrim et al. inoculated traditional yogurts containing *Streptococcus thermophilus* and *Lactobacillus bulgaricus* with *E. coli* O157:H7 and found that the pathogen was not detectable up to 10 days [27]. In addition, suppression of *S. aureus* might be due to its inability in catalase production to degrade H₂O₂ produced by the starters under acidic condition [28].

Table 1- Microbial count (CFU/g) of traditional yogurts

Sample	Coliform	<i>E. coli</i>	Mold	Yeast	<i>S. aureus</i>
1	<10	0	<100	>100	0
2	<10	0	<100	>100	0
3	<10	0	<100	>100	0
4	<10	0	>100	>100	0
5	<10	0	>100	>100	0
6	<10	0	<100	>100	0
7	<10	0	>100	>100	0
8	<10	0	>100	>100	0
9	<10	0	<100	>100	0
10	<10	0	<100	>100	0
11	<10	0	<100	>100	0
12	<10	0	<100	>100	0
13	<10	0	<100	>100	0
14	<10	0	<100	>100	0
15	<10	0	<100	>100	0
16	<10	0	<100	>100	0
17	<10	0	<100	>100	0
18	<10	0	<100	>100	0
19	<10	0	<100	>100	0
20	<10	0	<100	>100	0
21	<10	0	<100	>100	0
22	<10	0	<100	>100	0
23	<10	0	>100	>100	0

Our results showed high level of yeast in all samples (Table 1). In agreement, Guzeler et al. found non-significant difference in count of yeast/mold compared to lactic acid bacteria in

kefir yogurt [29]. In accordance to several studies that examined diversity of beneficial microorganisms including yeasts in traditional dairies from different regions of Iran [30-32], the high

yeast level in our samples was expected. Importantly, Faghihi Shahrestani et al. reported yeast species of traditional milk in Iran through which *Yarrowia lipolytica*, *Saccharomyces cerevisiae*, and *S. paradoxus* were detected [33] that none of them is pathogen. *Kluyveromyces*, another fungus that has been detected in yogurt, adopts with high and low temperatures of fermentation and storage, respectively, and considered as harmless organism for human [34]. Importantly, its metabolites such as rennet and lactase are listed in GRAS list of US FDA [35]. On the other hand, Afzali et al. found antagonistic activity of *L. brevis* isolated from traditional cheeses in Iran against pathogenic yeasts thanks to the organic acids produced by the bacteria at high concentration [36].

Considering the all results reported by other studies, because microbial load of traditional products is region-specific, a complementary evaluation is needed to determine the yeasts' type in our study. However, a total count up to 100 CFU/g of yeast-mold is acceptable according to Iranian regulation that is limited to mold in the products containing yeast as starter [37]. To the contrary, what is concern is that count of mold was more than acceptable level in approximately 22% (fine out of 23) of samples (Table 1). Dairy products may be source of aflatoxin contamination that mainly produced by *Aspergillus* species. Molds or their toxins enter to milk through consumption of contaminated silage by lactating animals [33]. Therefore, monitoring of animal farms to alleviate fungal contamination in further products is recommended.

3-2- Starch content

Starch may be added to dairy products to increase viscosity or mimic fat perception in consumers [38]. Starch granules consist of amylose and amylopectin in form of helices and branch, respectively. Several hydrophobic molecules such as fatty acids can enter to internal space of amylose [39]. Dispersion of lipid structures in starch molecules leads to a same sensory perception as pure fat that is important in preparation of dietary foods. In addition, water interaction with starch molecules through hydrogen bonds limits water mobility and increase viscosity of the mixture [40]. In this regard, Singh and Byars observed that addition of starch-lipid complex to yogurt up to 4% made a consistent texture without syneresis [41]. This was corroborated in the study of Lobato-Calleros et al. about studying the effect of starch on syneresis of yogurt so that phase separation was decreased as a result of H₂O molecules bounded to hydrophilic sites of starch [42]. Another study reported that addition of starch-lipid complex up to 2% induced a dense phase of casein within a reduced coagulation time. The authors showed that starch-lipid complex jointed to the casein aggregates and might act as nucleus of casein agglomeration in yogurt [43]. Similar result in viscosity enhancement and creaminess was observed by Ares et al. after addition of starch to stirred yogurt [44]. Although, starch addition is a technical point that promotes healthiness of low-fat products for obese people, it is not desirable by the consumers interested in intake of dairy fat especially from traditional products. Our investigation showed that majority of the samples were free of starch and just two out of the all contained high amount of starch (Figure 1).




Figure 1- Qualitative determination of starch in traditional yogurts

4. Conclusion

A large number of traditional foods are produced in Iran of which dairy products are popular because they are easily prepared and distributed within the country. However, the products' qualification is of concern for home-made products not underwent thermal or chemical processes. Although, acidic nature of fermented dairies is a good point in repression of spoilage bacteria, cross-contamination or viability of yeasts and molds that are relatively resistant to low pH should not be neglected. According to our results, there was no concern about coliform, *E. coli*, and *S. aureus* in the yogurts. In comparison, yeasts level was higher than acceptable limit in the all samples. Based on several reports of other scientists, Iranian traditional dairies are source of healthy microorganisms and the high yeast count would not be concerning. However, characterization of the yeasts in the samples by molecular approaches is undertaken. Furthermore, five yogurts were contaminated by molds, which should be ignored for consumption. More than 90% of the samples contained no or little starch. It shows the desirable raw materials and appropriate methods used in yogurt preparation. In general, there was no widespread concern about consumption of the traditional yogurts in north of Iran.

5. Conflict of interest

There is no conflict of interest to be declared.

References

- Mahan LK, Raymond JL. Krause's food and the nutrition care process. 14th editon, Elsevier; 2017.
- Pei R, Martin DA, DiMarco DM, Bolling BW. Evidence for the effects of yogurt on gut health and obesity. *Critical Review in Food Science and Nutrition*. 2017; 57(8): 1569-1583.
<https://doi.org/10.1080/10408398.2014.883356>
- Gomez-Gallego C, Gueimonde M, Salminen S. The role of yogurt in food-based dietary guidelines. *Nutrition Reviews*. 2018; 76(Supplement 1): 29-39.
<https://doi.org/10.1093/nutrit/nuy059>
- Shah NP. *Yogurt in health and disease prevention*. Academic Press; 2017.
- Tajabadi Ebrahimi M, Ouwehand AC, Hejazi MA, Jafari P. Traditional Iranian dairy products: a source of potential probiotic Lactobacilli. *African Journal of Microbiology Research*. 2011; 5(1): 20-27.

<https://doi.org/10.5897/AJMR10.629>

- Faghfoori Z, Gargari BP, Saber A, Seyyedi M, Khosroushahi AY. The investigation of the diversity of *Lactobacilli* spp. and assessment their some probiotic properties in traditional dairy products in east Azerbaijan province in Iran. *Iranian Journal of Pharmaceutical Research*. 2017; 16(4): 1538-1545.
- Sharifi Yazdi MK, Davoodabadi A, Khesht Zarin HR, Tajabadi Ebrahimi M, Soltan Dallal MM. Characterisation and probiotic potential of lactic acid bacteria isolated from Iranian traditional yogurts. *Italian Journal of Animal Science*. 2017; 16(2): 185-188.
<http://dx.doi.org/10.1080/1828051X.2016.1222888>
- Zarei F, Nateghi L, Eshaghi MR, Ebrahimi Tajabadi M. Optimization of gamma-aminobutyric acid production in probiotics extracted from local dairy products in west region of Iran using MRS broth and whey protein media. *Applied Food Biotechnology*. 2018; 5(4): 233-242.
<http://dx.doi.org/10.22037/afb.v5i4.22360>
- Hamzehlou P, Akhavan Sepahy A, Mehrabian S, Hosseini F. Production of vitamins B₃, B₆ and B₉ by Lactobacillus isolated from traditional yogurt samples from 3 cities in Iran, winter 2016. *Applied Food Biotechnology*. 2018; 5(2): 107-120.
<https://dx.doi.org/10.22037/afb.v5i2.18905>
- Vinceti M, Filippini T, Wise LA. Environmental selenium and human health: an update. *Current Environmental Health Reports*. 2018; 5(4): 464-485.
<https://doi.org/10.1007/s40572-018-0213-0>
- Shakibaie M, Mohammadi-Khorsand T, Adeli-Sardou M, Jafari M, Amirpour-Rostami S, Ameri A, Forootanfar H. Probiotic and antioxidant properties of selenium-enriched *Lactobacillus brevis* LSe isolated from an Iranian traditional dairy product. *Journal of Trace Elements in Medicine and Biolology*. 2017; 40: 1-9.
<https://doi.org/10.1016/j.jtemb.2016.11.013>
- Papadimitriou CG, Vafopoulou-Mastrojiannaki A, Silva SV, Gomes AM, Malcata FX, Alichanidis E. Identification of peptides in traditional and probiotic sheep milk yoghurt with angiotensin I-converting enzyme (ACE)-inhibitory activity. *Food Chemistry*. 2007; 105(2): 647-656.
<https://doi.org/10.1016/j.foodchem.2007.04.028>
- Dirisu C, Gloria L, Ebere I. Microbiological load of yoghurt sold in Omoku schools, Rivers State, Nigeria. *African Journal of Microbiology Research*. 2015; 9(34): 1960-1963.
<https://doi.org/10.5897/AJMR2015.7556>

14. International Standard Organization. Milk and milk products-general guidance for the preparation of test samples, initial suspensions and decimal dilutions for microbiological examination. ISO 8261. 2001.
15. International Standard Organization. Milk and milk products- enumeration of presumptive *Escherichia coli*- Part 1: Most Probable Number technique using 4-methylumbelliferyl- β -D-glucuronide (MUG). ISO 11866-1 IDF170-1. 2005.
16. International Standard Organization. Microbiology of food and animal feeding stuffs- Horizontal method for the enumeration of coagulase-positive staphylococci (*Staphylococcus aureus* and other species)-Part 1: Technique using Baird-Parker Agar medium. ISO 6888-1. 2003.
17. International Standard Organization. Microbiology of food and animal feeding stuffs- Horizontal method for the enumeration of yeasts and moulds- Part 1: Colony count technique in products with water activity greater than 0.95. ISO 21527-1. 2009.
18. Food Safety and Standards Authority of India. Manual of methods of analysis of foods, milk and milk products. 2015: pp. 9.
19. Codex Alimentarius. Codex standard for fermented milks: Codex Stan 243-2003. FAO United Nations: Roma. 2011.
20. Kaur M, Kaur J. Dairy products and their adulteration. Research & Reviews: Journal of Dairy Science and Technology. 2018; 7(2): 11-14. <https://doi.org/10.37591/rjodst.v7i2.869>
21. Xu L, Yan SM, Cai CB, Wang ZJ, Yu XP. The feasibility of using near-infrared spectroscopy and chemometrics for untargeted detection of protein adulteration in yogurt: removing unwanted variations in pure yogurt. Journal of Analytical Methods in Chemistry. 2013: 1-9. <https://doi.org/10.1155/2013/201873>
22. Azad T, Ahmed S. Common milk adulteration and their detection techniques. International Journal of Food Contamination. 2016; 3(1): 1-9. <https://doi.org/10.1186/s40550-016-0045-3>
23. Kim MJ, Kim HY. Direct duplex real-time loop mediated isothermal amplification assay for the simultaneous detection of cow and goat species origin of milk and yogurt products for field use. Food Chemistry. 2018; 246: 26-31. <https://doi.org/10.1016/j.foodchem.2017.11.014>
24. Abernethy, GA, Bendall JG, Holroyd SE. Advances in testing for adulteration and authenticity of dairy products. Advances in Food Authenticity Testing. 2016: 461-490. <https://doi.org/10.1016/B978-0-08-100220-9.00017-5>
25. Karacaglar NNY, Bulat T, Boyaci IH, Topcu A. Raman spectroscopy coupled with chemometric methods for the discrimination of foreign fats and oils in cream and yogurt. Journal of Food and Drug Analysis. 2019; 27(1): 101-110. <https://doi.org/10.1016/j.jfda.2018.06.008>
26. Russo P, Spano G, Capozzi V. Safety evaluation of starter cultures. In: Speranza B, Bevilacqua A, Corbo MR, Sinigaglia M. Starter cultures in food production. 1st edition. John Wiley & Sons. 2017: 101-128. <https://doi.org/10.1002/9781118933794.ch6>
27. Cutrim CS, deBarros RF, daCosta MP, Franco RM, Conte-Junior CA, Cortez MAS. Survival of *Escherichia coli* O157: H7 during manufacture and storage of traditional and low lactose yogurt. LWT. 2016; 70: 178-184. <https://doi.org/10.1016/j.lwt.2016.02.047>
28. Attaie R, Whalen PJ, Shahani KM, Amer MA. Inhibition of growth of *Staphylococcus aureus* during production of acidophilus yogurt. Journal of Food Protection. 1987; 50(3): 224-228. <https://doi.org/10.4315/0362-028X-50.3.224>
29. Guzeler N, Ari E, Konuray G, Ozbek C. Physicochemical and microbiological properties of kefir, kefir yogurt and chickpea yogurt. International Journal of Nutrition and Food Engineering. 2019; 13(7): 189-192.
30. Hajhosseinali M, Nasr S, Amoozegar MA, Yurkov A. *Saccharomycopsis oxydans* sp. nov., a new non-fermentative member in the genus *Saccharomycopsis* isolated from a traditional dairy product of Iran. International Journal of Systematic and Evolutionary Microbiology. 2019; 70(2): 1059-1063. <https://doi.org/10.1099/ijsem.0.003874>.
31. Karimpour F, Razavi SH, Nubar Tkhruni F. Investigation of antibacterial properties of yeast strains isolated from Iranian richal and traditional dairy products in Armenia. Armaghane Danesh. 2016; 21(6): 552-562.
32. Rahmati F. Characterization of *Lactobacillus*, *Bacillus* and *Saccharomyces* isolated from Iranian traditional dairy products for potential sources of starter cultures. AIMS Microbiology. 2017; 3(4): 815-825. <https://doi.org/10.3934/microbiol.2017.4.815>

33. Shahrestani FF, Tajabadi Ebrahimi M, Bayat M, Hashemi J, Razavilar V. The study of phenotypic and genotypic importance of fungal isolated from traditional milk and cheese in Iran. *Gene Reports*. 2019; 17: 1-4.
<https://doi.org/10.1016/j.genrep.2019.100472>
34. Moineau-Jean A, Guevremont E, Champagne CP, Roy D, Raymond Y, LaPointe G. Fate of *Escherichia coli* and *Kluyveromyces marxianus* contaminants during storage of Greek-style yogurt produced by centrifugation or ultrafiltration. *International Dairy Journal*. 2017; 72: 36-43.
<https://doi.org/10.1016/j.idairyj.2017.04.002>
35. US Food and Drug Administration. Microorganisms and microbial-derived ingredients used in food (Partial list). 2018. Available at:
<https://www.fda.gov/food/generally-recognized-safe-gras/microorganisms-microbial-derived-ingredients-used-food-partial-list>. 2018.
36. Afzali S, Edalatian Dovom MR, Habibi Najafi MB, Mazaheri Tehrani M. Determination of the anti-yeast activity of *Lactobacillus* spp. isolated from traditional Iranian cheeses in vitro and in yogurt drink (Doogh). *Scientific Reports*. 2020; 10(1): 1-11.
37. Iranian National Standard Organization. Microbiology of milk and milk products-Specifications and test methods. 3rd Revision. INSO 2406. 2017.
38. Radi M, Niakousari M, Amiri S. Physicochemical, textural and sensory properties of low-fat yogurt produced by using modified wheat starch as a fat replacer. *Journal of Applied Sciences*. 2009; 9(11): 2194-2197.
39. Lu X, Shi C, Zhu J, Li Y, Huang Q. Structure of starch-fatty acid complexes produced via hydrothermal treatment. *Food Hydrocolloids*. 2019; 88: 58-67.
<https://doi.org/10.1016/j.foodhyd.2018.09.034>
40. Luchese CL, Spada JC, Tessaro IC. Starch content affects physicochemical properties of corn and cassava starch-based films. *Industrial Crops and Products*. 2017; 109: 619-626.
<https://doi.org/10.1016/j.indcrop.2017.09.020>
41. Singh M, Byars JA. Starch-lipid composites in plain set yogurt. *International Journal of Food Science and Technology*. 2009; 44(1): 106-110.
<https://doi.org/10.1111/j.1365-2621.2007.01661.x>
42. Lobato-Calleros C, Ramirez-Santiago C, Vernon-Carter E, Alvarez-Ramirez J. Impact of native and chemically modified starches addition as fat replacers in the viscoelasticity of reduced-fat stirred yogurt. *Journal of Food Engineering*. 2014; 131: 110-115.
<https://doi.org/10.1016/j.jfoodeng.2014.01.019>
43. Singh M, Kim S. Yogurt fermentation in the presence of starch-lipid composite. *Journal of Food Science*. 2009; 74(2): 85-89.
<https://doi.org/10.1111/j.1750-3841.2008.01028.x>
44. Ares, G., Perez C, Reolon G, Segura N, Lema P, Gambaro A. Influence of gelatin and starch on the instrumental and sensory texture of stirred yogurt. *International Journal of Dairy Technology*. 2007; 60(4): 263-269.
<https://doi.org/10.1111/j.1471-0307.2007.00346.x>